The Universe as a Quantum Computer: Qubit Distribution from CMB Photons to Supermassive Black Holes

Douglas C. Youvan

doug@youvan.com

www.youvan.ai

May 22, 2026

The conceptualization of the universe as a vast quantum computational system has fundamentally altered the intersection of quantum information theory, astrophysics, and cosmology. Rather than treating information as a mere mathematical abstraction, modern physics increasingly recognizes it as a foundational physical property inherently tied to matter, energy, and entropy. This paradigm shift naturally provokes a profound cosmological question regarding the physical location and state of the most abundant qubits in the universe. The answer necessitates a rigorous distinction between the sheer volume of free-floating, non-interacting quantum states and the concentrated density of gravitationally bound information. This paper comprehensively explores the cosmological distribution of quantum information by identifying its primary repositories. First, we examine the Cosmic Microwave Background and Cosmic Neutrino Background, which permeate the interstellar vacuum and account for the absolute greatest number of discrete qubits. Second, we analyze supermassive black holes through the lens of the holographic principle and Bekenstein-Hawking entropy, establishing them as the densest localized concentrations of quantum information. Finally, we evaluate interstellar hydrogen clouds as the most abundant atomic-scale qubits. By mapping these cosmic data structures, this paper constructs a theoretical framework for calculating universal computational capacity from the Big Bang to the present epoch.

Keywords: quantum information, cosmology, qubits, cosmic microwave background, supermassive black holes, Bekenstein-Hawking entropy, holographic principle, universal computation, cosmic neutrino background, quantum astrophysics.

A collaboration with Google Gemini-3.1-Pro. CC4.0.

Outline

1. Introduction to Quantum Information Cosmology 1.1 The Information-Theoretic Universe Paradigm 1.2 Historical Context of Universal Computation 1.3 Scope and Methodology of the Paper

2. Defining the Cosmological Qubit 2.1 Quantum Degrees of Freedom in Fundamental Particles 2.2 Photonic, Fermionic, and Atomic Two-State Systems 2.3 Distinguishing Between Free and Bound Information

3. The Computational Capacity and Limits of the Cosmos 3.1 Theoretical Bounds on Universal Processing Power 3.2 Calculating Total Information Content from the Big Bang to Present 3.3 The Lloyd Limit and Physical Constraints on Computation

4. The Cosmic Microwave Background: An Ocean of Photonic Qubits 4.1 Origin and Density of CMB Photons 4.2 Polarization and Spin States as Information Carriers 4.3 The Total Qubit Volume of the Observable CMB

5. The Cosmic Neutrino Background: Elusive Quantum States 5.1 Relic Neutrinos and Weak Interactivity 5.2 Flavor Oscillations and Superposition 5.3 Comparing Neutrino Abundance to the CMB

6. Baryonic Qubits and the Interstellar Medium 6.1 Hydrogen as the Universe's Default Atomic Qubit 6.2 Spin-Flip Transitions and the 21-centimeter Line 6.3 Nebular Hydrogen Clouds as Dilute Quantum Processors

7. Black Hole Thermodynamics and Entropy 7.1 The Event Horizon and the Area Law 7.2 Deriving Bekenstein-Hawking Entropy 7.3 Translating Entropy into Qubits of Hidden Information

8. Supermassive Black Holes: The Densest Information Repositories 8.1 Galactic Centers and Extreme Mass Concentrations 8.2 Information Density Comparisons: Vacuum vs. Singularity 8.3 The Black Hole Information Paradox in a Computational Context

9. The Holographic Principle and Universal Boundaries 9.1 Encoding 3D Volume on a 2D Boundary 9.2 Maximum Entropy Bounds for the Observable Universe 9.3 Implications for the Spatial Distribution of Qubits

10. Dark Matter and Dark Energy: Unknowns in Cosmic Computation 10.1 Potential Information Carrying Capacity of Dark Matter Candidates 10.2 Dark Energy and the Expanding Computational Space 10.3 Theoretical Models for Dark Qubits

11. Cosmological Quantum Entanglement 11.1 Primordial Entanglement and Cosmic Inflation 11.2 Bell Correlations across the Cosmic Microwave Background 11.3 Decoherence and the Classical Transition of the Universe

12. Future Prospects in Observational Quantum Cosmology 12.1 Proposed Detection Methods for Cosmological Qubit Signatures 12.2 Integrating Quantum Computing with Astrophysical Data 12.3 Final Conclusions on the Universe's Computational Architecture

13. References

Art

1. Introduction to Quantum Information Cosmology

The intersection of cosmology and quantum information theory has generated one of the most profound conceptual shifts in modern physics. Traditionally, cosmology has sought to understand the universe through the lenses of classical mechanics, general relativity, and quantum field theory, focusing heavily on the physical interactions of fundamental forces and matter over vast expanses of space and time. However, the emergence of quantum information cosmology introduces a radically different fundamental currency to describe reality: the qubit. In this theoretical framework, the universe is not merely an inert stage where physical processes occur, but rather a dynamic, evolving computational network. Every subatomic interaction constitutes a quantum logic gate, and every elementary particle serves as a physical register of information. This introductory section establishes the foundational concepts of treating the cosmos as an information-processing system, contextualizes this paradigm within the historical evolution of theoretical physics, and outlines the structural methodology used throughout this paper to map the distribution of cosmic qubits from the Big Bang to the present epoch.

1.1 The Information-Theoretic Universe Paradigm

The information-theoretic view of the universe posits that physical reality is fundamentally built upon information rather than just matter and energy. This represents a significant departure from the historical materialistic view where particles and continuous fields are considered the ultimate bedrock of reality. Instead, the current paradigm suggests that the complete physical state of any system can and must be described by the quantum information encoded within it. This perspective is anchored in the realization that information is not an abstract mathematical concept invented by human observers, but a tangible physical entity subjected strictly to the laws of thermodynamics and quantum mechanics. The erasure of information inevitably dissipates heat, proving that computational states are irrevocably tied to physical states.

When a cosmic microwave background photon travels through the vacuum of space, or when electrons and protons interact in a primordial nebula to form neutral hydrogen, they are effectively processing, storing, and transmitting quantum information. A quantum bit, or qubit, differs radically from a classical bit in that it can exist in a complex superposition of states, allowing for the massive parallel processing capacity that governs quantum mechanics. By mapping the intrinsic degrees of freedom of fundamental particles, such as particle spin, photon polarization, and spatial coordinates, to the discrete states of a theoretical qubit, physicists can calculate the absolute informational capacity of any region of space. Consequently, the expansion of the universe, the gravitational formation of massive cosmic structures, and the overall thermodynamic evolution of the cosmos can all be seamlessly reinterpreted as the ongoing execution of a universal quantum algorithm.

1.2 Historical Context of Universal Computation

The conceptual lineage of the universe as a computer traces its roots through several key, intersecting developments in physics, thermodynamics, and computer science during the twentieth century. The theoretical foundation was initially laid by early pioneers of classical computation, who established that any logical physical process could theoretically be simulated by a universal computing machine. However, as the bizarre rules of quantum mechanics matured and became better understood, it became mathematically evident that classical computation could never efficiently simulate complex quantum phenomena. This friction led to the eventual proposition that fundamental computational systems must inherently operate on quantum principles if they are to accurately reflect nature.

Later in the twentieth century, the phrase "it from bit" was introduced to the physics lexicon, encapsulating the radical idea that all physical entities are ultimately derived from fundamentally information-theoretic origins, requiring a series of binary quantum queries to define their reality. Simultaneously, the groundbreaking study of black hole thermodynamics revealed a deep, unexpected connection between gravity, thermodynamic entropy, and quantum information. The mathematical discovery that the entropy of a black hole is directly proportional to the surface area of its event horizon, rather than its three-dimensional volume, demonstrated a strict physical limit to how densely information can be packed into a region of space. This realization subsequently birthed the holographic principle. In the early twenty-first century, these varied theoretical threads were synthesized into formal computational models capable of calculating the maximum processing capacity of the universe, setting hard physical bounds on the total number of logical operations that could have possibly occurred since the Big Bang. These historical milestones successfully transitioned the idea of cosmic computation from an intriguing philosophical metaphor into a rigorously quantifiable branch of modern astrophysics.

1.3 Scope and Methodology of the Paper

This paper aims to systematically categorize, locate, and quantify the primary repositories of quantum information distributed across the observable universe. To achieve this comprehensive overview, the paper is structured to analyze both the most ubiquitous free-floating fundamental particles and the most extreme localized concentrations of gravitationally bound mass-energy. The scope primarily encompasses the relic radiation leftover from the Big Bang, specifically the Cosmic Microwave Background and the Cosmic Neutrino Background, which represent the absolute highest raw count of distributed, non-interacting qubits in existence. The analysis then shifts to baryonic matter, focusing heavily on interstellar hydrogen clouds as the primary atomic-scale qubit repository. Subsequently, the paper examines the severe gravitational extremes of

supermassive black holes, utilizing established thermodynamic entropy formulas to quantify their unparalleled and massive information density.

The methodology of this paper relies on synthesizing established cosmological observational data, including particle number densities, cosmic expansion rates, and galactic mass distributions, with the rigid mathematical principles of quantum information theory. By strictly applying theoretical limits of physical computation to astrophysical observables, this paper constructs a comprehensive and cohesive map of the universe's informational architecture. No novel observational data is presented; rather, existing astrophysical metrics are translated into information-theoretic values using standard conversion formulas. The following sections will sequentially expand upon these specific cosmic data repositories, beginning with a strict physics-based definition of the cosmological qubit and the fundamental mathematical limits of universal processing power.

2. Defining the Cosmological Qubit

To map the computational capacity of the universe, one must first establish a rigorous definition of what constitutes a cosmological qubit. In human-engineered quantum computers, qubits are artificially constructed using delicate macroscopic systems such as superconducting circuits, trapped ions, or precisely controlled neutral atoms. These laboratory setups are designed to isolate a two-state quantum system from the chaotic external environment to prevent sudden decoherence. However, when observing the universe at large, nature provides its own fundamental quantum information registers. In the cosmological context, a qubit is not a manufactured device but an intrinsic property of matter and energy. It represents the smallest discrete unit of quantum information naturally occurring in the universe. Understanding how this information is physically instantiated requires breaking down fundamental particles into their measurable properties and identifying the specific quantum states that can act as binary variables. This section defines the physical parameters that allow elementary particles to function as qubits, categorizes the primary types of naturally occurring two-state systems, and establishes a critical theoretical distinction between information that travels freely through the cosmic void and information that is permanently bound by gravitational and nuclear forces.

2.1 Quantum Degrees of Freedom in Fundamental Particles

The storage capacity of any physical system, whether it is a classical hard drive or a quantum universe, is strictly dictated by its available degrees of freedom. In physics, a degree of freedom refers to an independent physical parameter that is required to fully define the state of a system. For fundamental particles, these parameters include spatial coordinates, momentum vectors, electrical charge, and intrinsic angular momentum, commonly known as spin. Classical

mechanics historically assumed these properties could take on a continuous, infinite range of values, which would imply that even a single particle could theoretically store an infinite amount of data. However, the fundamental laws of quantum mechanics forbid this. The quantization of space and energy, coupled with the Heisenberg uncertainty principle, divides the phase space of any particle into discrete, finite cells.

Because a particle can only occupy a finite number of distinguishable states, its capacity to hold information is strictly bounded. The most fundamental building block of this information is the qubit, which corresponds to a quantum system that possesses exactly two mutually exclusive, observable states. While a specific particle may have multiple degrees of freedom, any isolated property that can be measured as one of two possible outcomes effectively acts as a solitary qubit. The immense computational power of the universe does not stem from an infinite storage capacity within individual subatomic particles, but rather from the sheer abundance of these discrete degrees of freedom spread across the cosmos. When these fundamental particles interact through the fundamental forces of nature, their respective degrees of freedom become entangled, essentially executing complex logical operations that mathematically drive the physical evolution of the universe.

2.2 Photonic, Fermionic, and Atomic Two-State Systems

To quantify the universe's qubits, we must identify the specific physical phenomena that naturally form two-state quantum systems. These systems can be broadly categorized into photonic, fermionic, and atomic structures. Photons, the gauge bosons of the electromagnetic force, are perhaps the most ubiquitous carriers of quantum information. The primary qubit encoded within a photon is its polarization. A single photon can be polarized horizontally or vertically, or in a quantum superposition of both, perfectly mirroring the zero and one states of a computational qubit. Because photons travel at the speed of light and rarely interact with one another in a vacuum, their polarization states are excellent preservers of pristine quantum information over vast cosmological distances.

Fermions, the fundamental building blocks of ordinary matter such as electrons, protons, and neutrinos, encode information primarily through their intrinsic spin. A fermion possesses a half-integer spin that can be measured as either spin-up or spin-down relative to a specific observational axis. This binary orientation provides a natural and highly robust qubit. Finally, when fermions bind together to form atoms, new composite quantum states emerge. The simplest atomic two-state system is found in the hydrogen atom, where the relative spin alignments of its solitary electron and proton create distinct, measurable energy levels. The transition between the parallel and antiparallel spin states of neutral hydrogen represents a pure atomic-scale qubit. Together, these photonic polarizations, fermionic spins, and atomic

energy levels constitute the foundational hardware architecture of the cosmic quantum computer.

2.3 Distinguishing Between Free and Bound Information

Having established the physical nature of cosmological qubits, it is necessary to classify how this information is distributed and processed across spatial dimensions. In the information-theoretic paradigm, cosmic data can be broadly divided into two distinct categories: free information and bound information. Free information is characterized by fundamental particles that travel essentially unimpeded through the vacuum of space. The most prominent examples are the low-energy photons of the Cosmic Microwave Background and the highly elusive particles of the Cosmic Neutrino Background. These free qubits are incredibly dilute in terms of spatial density, yet they represent the largest aggregate sum of discrete information in the observable universe due to their staggering total numbers. They function as the universe's long-term memory, carrying primordial data from the very early stages of the Big Bang with minimal computational disruption, interaction, or decoherence.

Conversely, bound information refers to qubits that are trapped within complex, interacting, and gravitationally confined structures. This encompasses everything from dense interstellar gas clouds and stellar interiors to terrestrial planets and, ultimately, the extreme environments of black holes. In these localized astrophysical systems, particles continuously collide and interact, causing their quantum states to rapidly entangle, compute, and decohere. Bound information acts as the active processing unit of the universe, where the thermodynamic state of matter is constantly being dynamically updated. While bound qubits are numerically fewer than their free-floating counterparts across the totality of the cosmos, their immense spatial concentration leads to extreme localized information densities. This crucial distinction between the dispersed, static memory of free cosmic radiation and the localized, active processing of bound mass-energy dictates how quantum information behaves on a macroscopic scale and forms the structural basis for analyzing the cosmos as a cohesive computational entity.

3. The Computational Capacity and Limits of the Cosmos

To treat the universe as a quantum computer requires more than just identifying the physical nature of its qubits; it demands a rigorous quantification of its ultimate processing power. Just as a classical computer is constrained by the physical limitations of its hardware—such as processor clock speeds, memory limits, and thermal cooling requirements—the universe itself is strictly bounded by the fundamental laws of physics. Thermodynamics, quantum mechanics, and general relativity impose inescapable mathematical ceilings on how much information a given region of space can hold and how fast that information can be processed. By applying

these physical limits to the observable mass and energy of the cosmos, physicists can calculate the absolute maximum computational capacity of the universe. This section explores the fundamental theorems that dictate physical computation rates, calculates the total quantum information content available since the Big Bang, and examines the hard constraints that govern the universe's overall informational evolution.

3.1 Theoretical Bounds on Universal Processing Power

The ultimate speed at which any physical system can process information is governed by the Margolus-Levitin theorem. Formulated in the late twentieth century, this theorem bridges quantum mechanics and computer science by establishing that the maximum rate at which a quantum system can transition from one state to an orthogonal, perfectly distinguishable state is strictly proportional to the system's average energy. Specifically, the maximum number of logical operations per second is equal to twice the energy of the system divided by Planck's constant times pi. Because energy is equivalent to mass according to special relativity, the processing speed of any physical computing device is ultimately bottlenecked by the amount of mass-energy it contains.

A parallel constraint is known as Bremermann's limit, which calculates the maximum computational speed of a self-contained system based on its mass, utilizing the speed of light and Planck's constant. Bremermann deduced that one gram of matter can execute a maximum of roughly 1.36 x 10^47 logical operations per second. When these quantum bounds are applied to the macroscopic universe, the entire mass-energy of the observable cosmos becomes the theoretical power supply for a single, interconnected processing unit. If every ounce of matter and radiation in the universe were perfectly harnessed to perform quantum operations, the laws of physics dictate a hard ceiling on the speed of cosmic computation. These physical bounds prove that the universe is not processing an infinite amount of data at infinite speeds; rather, it is a finite system governed by a precise and measurable thermodynamic clock.

3.2 Calculating Total Information Content from the Big Bang to Present

To determine the sheer size of the cosmic quantum computer's memory, one must look at the thermodynamic entropy of the universe, which serves as a direct proxy for its total information content. Entropy is a measure of the number of microscopic arrangements that correspond to a specific macroscopic state. In information theory, this is mathematically identical to counting the number of physical bits or qubits required to perfectly describe the system.

By analyzing the current composition of the observable universe, physicists can calculate its total memory capacity. The vast majority of standard computational memory resides in the background radiation left over from the Big Bang. When the total entropy of the Cosmic Microwave Background and the Cosmic Neutrino Background is calculated, it yields roughly

10^90 bits of information. However, this number only accounts for free-floating radiation. When the extreme entropy of supermassive black holes is factored into the equation, the total informational content of the observable universe jumps by several orders of magnitude, reaching approximately 10^104 bits.

Furthermore, if one applies the holographic principle to the cosmological event horizon—the absolute boundary of the observable universe determined by the speed of light and cosmic expansion—the theoretical limit of total cosmic information reaches an astounding 10^120 bits. To measure the extent of the universe's computation over its 13.8 billion-year history, physicists multiply the total available mass-energy by the age of the universe. This calculation reveals that since the Big Bang, the universe could have performed a maximum of roughly 10^120 elementary quantum logical operations on 10^90 to 10^120 discrete qubits.

3.3 The Lloyd Limit and Physical Constraints on Computation

The synthesis of these cosmic limits is most famously formalized in the Lloyd limit, named after the physicist who first calculated the ultimate computational capacity of the universe. The Lloyd limit serves as the absolute boundary for any computation occurring within our physical reality. It demonstrates that the speed of light limits how fast information can travel from one qubit to another, Planck's constant limits how fast those qubits can flip between distinct states, and the gravitational constant limits how much information can be packed into a finite space before collapsing into a black hole.

This framework imposes strict physical constraints on the nature of the cosmic computer. Because the universe has a finite age and a finite speed of light, there are absolute horizons that isolate different regions of the cosmic processor from one another. Distant regions of the universe cannot instantly exchange information, meaning the cosmic computer is not a single, highly integrated processor, but rather a massively parallel, highly fragmented distributed computing network. Information is locally processed at maximum efficiency within gravitationally bound systems like galaxies and star clusters, while data transfer between these massive nodes occurs via the incredibly slow transit of photons across millions of light-years of empty space. The Lloyd limit ultimately proves that while the universe possesses an unimaginably large computational capacity, it is inherently restricted by the very physical laws it computes, forever preventing instantaneous, universe-wide quantum entanglement or infinite data processing.

4. The Cosmic Microwave Background: An Ocean of Photonic Qubits

When transitioning from the theoretical limits of cosmic computation to the physical location of the universe's data, the most obvious and pervasive structure to analyze is the Cosmic

Microwave Background. If the universe is a quantum computer, the Cosmic Microwave Background represents its largest and most enduring physical memory bank. Permeating every cubic centimeter of the observable cosmos, this relic radiation constitutes the sheer numerical majority of all free-floating fundamental particles in existence. To understand the universe's baseline computational capacity, one must view these ancient photons not merely as thermal remnants of a primordial explosion, but as individual, discrete physical registers carrying primordial quantum information. This section explores the cosmological origins and sheer physical density of these photons, details the exact quantum mechanics that allow them to function as individual photonic qubits, and calculates the staggering aggregate volume of quantum data continuously washing across the cosmic vacuum.

4.1 Origin and Density of CMB Photons

The origin of this vast photonic data ocean dates back to a pivotal moment in cosmological history known as the epoch of recombination, which occurred approximately 380,000 years after the initial Big Bang. Prior to this epoch, the universe was profoundly hot and dense, consisting of an opaque, churning plasma of free electrons and protons. In this state, photons could not travel any significant distance before violently colliding with free electrons through a process known as Thomson scattering. The early universe was essentially a closed thermodynamic system where quantum information was constantly being scrambled, entangled, and violently decohered at a massive rate.

However, as the universe expanded, its ambient temperature naturally plummeted. When the temperature dropped to approximately 3,000 Kelvin, the thermodynamic energy of the plasma was no longer sufficient to keep electrons completely severed from protons. In a sweeping, universe-wide phase transition, electrons rapidly bound to protons to form the first stable, electrically neutral hydrogen and helium atoms. Because neutral atoms do not scatter light in the same way free electrons do, the universe suddenly transformed from an opaque, dense fog into a vast, transparent vacuum. The photons that existed at that exact moment were abruptly freed from their chaotic interactions. They decoupled from matter entirely and began streaming freely across the expanding cosmos.

Today, these decoupled photons are observed as the Cosmic Microwave Background. Over the past 13.8 billion years, the relentless expansion of space has stretched their wavelengths, shifting them out of the visible spectrum and into the microwave frequency, while dropping their temperature to a frigid 2.7 Kelvin. Despite this extreme loss of energy, the photons themselves remain fully intact. Modern cosmological measurements, most notably those taken by satellites mapping the cosmic sky, reveal an astonishing density: there are approximately 411 Cosmic Microwave Background photons in every single cubic centimeter of empty space. Because the vacuum of space is so profoundly vast, this relatively modest local density

accumulates into an unimaginably massive total population, making these photons the most abundant identifiable entities in the physical universe.

4.2 Polarization and Spin States as Information Carriers

Identifying the sheer number of photons is only the first step in mapping the cosmic computer; one must also understand exactly how a single, free-floating photon physically encodes quantum data. A photon is an elementary particle classified as a gauge boson, mediating the electromagnetic force. Because a photon is entirely massless and travels exactly at the speed of light, the laws of quantum mechanics restrict its available degrees of freedom compared to particles with mass. Specifically, a photon possesses an intrinsic angular momentum, or spin, of 1. However, due to its massless nature, the spin vector can only point strictly parallel or strictly anti-parallel to its direction of motion.

These two distinct states are known as helicity, manifesting physically as either right-handed or left-handed circular polarization. Because there are exactly two mutually exclusive, orthogonal states available to the photon's spin, a single photon acts as a perfect, naturally occurring two-level quantum system. It is a pristine physical qubit. The state of this photonic qubit can be a discrete zero, a discrete one, or, crucially, a complex quantum superposition of both linear and circular polarization states.

What makes the Cosmic Microwave Background so uniquely vital to the concept of universal memory is the exceptionally high fidelity of its qubits. In human-engineered quantum computers, the greatest obstacle is decoherence, where a qubit loses its quantum state due to interacting with the surrounding environment. The vacuum of space, however, is a near-perfect isolator. Once the primordial photons decoupled from matter during the epoch of recombination, the vast majority of them never interacted with another particle again. They have been traveling in a straight line through an expanding vacuum for nearly 14 billion years. Consequently, the polarization state of a Cosmic Microwave Background photon today is a perfectly preserved, unaltered piece of quantum information directly inherited from the thermodynamic conditions of the early universe. They are the ultimate long-term data storage mechanism, immune to degradation, passively carrying the initial boundary conditions of the cosmos across the expanse of time.

4.3 The Total Qubit Volume of the Observable CMB

To comprehend the scale of this data repository, one must calculate the absolute aggregate volume of these photonic qubits across the observable universe. The observable universe is currently estimated to be a sphere with a comoving radius of approximately 46.5 billion light-years. Converting this immense scale into standard volume yields roughly 3.5 x 10^80 cubic meters, or 3.5 x 10^86 cubic centimeters.

By multiplying this total spatial volume by the measured density of 411 photons per cubic centimeter, physicists arrive at the total number of Cosmic Microwave Background photons currently residing within our observable horizon. The result is approximately 1.4 x 10^89 individual photons. Because each of these photons carries at least one discrete qubit of information via its polarization state, the Cosmic Microwave Background alone accounts for roughly 10^89 to 10^90 qubits of free-floating quantum information.

From a thermodynamic perspective, this overwhelming abundance of photons represents the dominant portion of the universe's baseline entropy, excluding the localized extremes of black holes. The radiation field of the Cosmic Microwave Background exists in a state of maximum thermal equilibrium, meaning its physical arrangement resembles random thermal noise. In information theory, maximum entropy correlates to the maximum amount of required information to describe a system. Therefore, while this vast ocean of 10^90 qubits may not be executing structured, purposeful algorithms like a biological brain or a silicon processor, it undeniably constitutes the sheer bulk of the universe's distributed physical memory. It is a ubiquitous, background hum of raw quantum data that dwarfs the information content of all the stars, planets, and galaxies combined.

5. The Cosmic Neutrino Background: Elusive Quantum States

While the Cosmic Microwave Background represents the most easily observable and numerically superior ocean of quantum information, it is accompanied by a slightly more elusive, yet equally fundamental, cosmic data repository. The Cosmic Neutrino Background consists of relic neutrinos that permeate the universe much like the primordial photons, yet they carry information from a significantly earlier and more violent epoch of cosmic history. Neutrinos are fundamentally different from photons in their particle nature, their interactions with other matter, and their intrinsic quantum behaviors. In the context of quantum information cosmology, these ghostly particles represent a massively parallel array of fermionic qubits executing a continuous, spontaneous quantum algorithm across billions of light-years. This section explores the primordial origins of relic neutrinos, examines their unique quantum property of macroscopic superposition through flavor oscillation, and calculates their total informational contribution to the cosmic computational network relative to the cosmic microwave photons.

5.1 Relic Neutrinos and Weak Interactivity

To understand the informational value of relic neutrinos, one must look back to the very first moments of the universe. The photons of the Cosmic Microwave Background decoupled from matter approximately 380,000 years after the Big Bang, which, in cosmological terms, is a

relatively late stage of early development. Neutrinos, however, decoupled drastically earlier. Approximately one second after the initial Big Bang, the universe was an incredibly dense, seething plasma of protons, neutrons, electrons, positrons, and neutrinos, with a temperature hovering around ten billion Kelvin. At this extreme temperature, neutrinos interacted constantly with other particles through the weak nuclear force.

However, as the universe rapidly expanded and the temperature dropped below one mega-electron volt, the density of the plasma decreased to the point where the weak nuclear force could no longer maintain thermal equilibrium between the neutrinos and the rest of the matter. In a fraction of a second, the neutrinos decoupled. From that moment forward, roughly one second into the universe's existence, these relic neutrinos have traveled freely through the expanding cosmos. Because they only interact via the weak nuclear force and gravity, they possess an infinitesimal cross-section for interaction. They can pass through entire planets, stars, and galaxies without ever scattering or collapsing their quantum states. From a quantum computing perspective, this extreme isolation makes relic neutrinos the most perfectly insulated, undisturbed qubits in the universe. They hold the deepest, oldest unaltered quantum records in existence, carrying raw data directly from the first second of time, entirely shielded from the decoherence that plagues baryonic matter.

5.2 Flavor Oscillations and Superposition

The most profound distinction between a photonic qubit and a neutrino qubit lies in how their quantum states evolve over time. Unlike photons, which are strictly massless and do not experience time, neutrinos possess a vanishingly small, yet non-zero, mass. This slight mass requires them to travel fractionally slower than the speed of light, meaning they possess an internal clock and experience the flow of time. In the Standard Model of particle physics, neutrinos come in three distinct flavors: electron, muon, and tau. However, a neutrino's flavor state does not perfectly align with its mass state. When a neutrino is born or interacts, it is in a definite flavor state, but this flavor state is actually a quantum superposition of three distinct mass states.

Because each of the three mass states propagates through the vacuum of space at slightly different quantum mechanical frequencies, the wave functions of the mass states drift out of phase with one another as the neutrino travels. This shifting phase relationship causes the probability amplitude of the overall flavor state to change. Consequently, a relic neutrino that began its journey as an electron neutrino billions of years ago may spontaneously be measured as a muon or tau neutrino today. This phenomenon, known as neutrino oscillation, is one of the only macroscopic, naturally occurring demonstrations of pure quantum superposition and interference. In the language of quantum information theory, a relic neutrino is not merely a static memory drive like a cosmic photon. Instead, its internal state is constantly undergoing a

unitary transformation. Every single free-floating neutrino in the universe is essentially a solitary quantum logic gate, continuously executing a complex phase-rotation algorithm across intergalactic distances.

5.3 Comparing Neutrino Abundance to the CMB

To map the full architecture of the universe's free-floating memory, the abundance of these fermionic qubits must be quantified and compared to the photonic qubits of the Cosmic Microwave Background. Following their decoupling one second after the Big Bang, the expansion of the universe stretched and cooled the relic neutrinos, analogous to the cosmic photons. Today, the Cosmic Neutrino Background exists at a staggering temperature of approximately 1.95 Kelvin, slightly colder than the microwave background.

Current cosmological models dictate the precise ratio of relic neutrinos to relic photons. For every cubic centimeter of the observable universe, there are approximately 339 relic neutrinos distributed across all three flavor states, equating to roughly 113 neutrinos per flavor. When multiplied by the sheer volume of the observable universe, this yields an aggregate total of roughly 1.2 x 10^89 free-floating relic neutrinos. While this total falls slightly short of the 1.4 x 10^89 photons in the microwave background, it still represents a profoundly massive repository of quantum information, firmly cementing neutrinos as the second most abundant discrete particles in the cosmos.

Furthermore, the informational characteristics of these two backgrounds differ significantly due to quantum statistics. Photons are bosons and are governed by Bose-Einstein statistics, meaning limitless numbers of them can occupy the exact same quantum state, which inherently limits the diversity of the information they carry. Neutrinos, on the other hand, are fermions, governed by Fermi-Dirac statistics and the Pauli exclusion principle. No two neutrinos can occupy the exact same quantum state in the same space. This restriction naturally forces the Cosmic Neutrino Background into a more highly structured, complex informational arrangement. Therefore, while relic neutrinos are marginally fewer in absolute number than cosmic photons, their fermionic nature, coupled with their continuous quantum flavor oscillations, makes them arguably the most complex and robust distributed quantum processing network traversing the cosmic vacuum.

6. Baryonic Qubits and the Interstellar Medium

Having mapped the vast, distributed networks of free-floating photons and relic neutrinos, the architectural analysis of the cosmic computer must now transition from the vacuum of space to the realm of structured mass. While the background radiation of the universe acts as an isolated, static memory drive, baryonic matter—the protons, neutrons, and electrons that

comprise the physical structures of the cosmos—represents the active, interactive processing components of the universal quantum computer. Unlike relic radiation, baryonic particles interact readily through all four fundamental forces of nature. These interactions allow their quantum states to become heavily entangled, driving the chemical and thermodynamic evolution of the cosmos. To understand how the universe computes complex structures like stars and galaxies, one must examine the simplest, most abundant building block of baryonic matter. This section identifies the neutral hydrogen atom as the universe's foundational atomic qubit, explores the specific quantum mechanics of its spin-flip transition as a localized logic operation, and evaluates the massive interstellar hydrogen clouds as dilute but highly active cosmic data processing centers.

6.1 Hydrogen as the Universe's Default Atomic Qubit

In the construction of artificial quantum computers, engineers often rely on trapping individual neutral atoms using optical tweezers to create highly stable, controllable qubits. Nature, in its cosmic architecture, utilizes the exact same principle but on a profoundly larger scale. The foundational hardware of the baryonic universe is the neutral hydrogen atom. Accounting for roughly 75 percent of all normal matter by mass, hydrogen is the simplest and most ubiquitous composite quantum system in existence. Consisting of a single proton nucleus orbited by a single electron, hydrogen provides an incredibly pristine and mathematically elegant quantum environment.

To function as a qubit, a system must possess two distinct, measurable quantum states. In the hydrogen atom, this two-state system is naturally provided by the intrinsic angular momentum, or spin, of its constituent particles. Both the proton and the electron are fermions with a spin of 1/2. Because they are bound together by the electromagnetic force, their magnetic dipole moments interact with one another. This interaction dictates that the overall ground state of the hydrogen atom is split into two distinct, closely spaced energy levels depending entirely on the relative orientation of the particle spins. When the spin of the electron is parallel to the spin of the proton, the atom is in a slightly higher energy state. When the spins are antiparallel, pointing in opposite directions, the atom rests in its absolute lowest energy state. The strict dichotomy between these parallel and antiparallel spin alignments creates a perfect, naturally occurring physical manifestation of a binary zero and one. Because hydrogen is overwhelmingly the most common element, these atomic-scale two-state systems are by far the most abundant localized memory registers of ordinary matter in the universe.

6.2 Spin-Flip Transitions and the 21-centimeter Line

The true computational value of a qubit lies not just in its ability to hold a state, but in its ability to undergo a state change, effectively executing a logical operation. For the hydrogen qubit, this

operation occurs via a specific quantum mechanical process known as a spin-flip transition. When a neutral hydrogen atom drops from the higher-energy parallel state to the lower-energy antiparallel state, it must shed the tiny difference in energy between the two configurations. It achieves this by emitting a single, highly specific photon.

Because the energy difference between these two magnetic spin states is incredibly minute—roughly 5.87 x 10^-6 electron volts—the emitted photon possesses a very low frequency of exactly 1420.4 megahertz. In the spatial domain, this corresponds to a wavelength of exactly 21.106 centimeters. This highly specific emission is known in astrophysics as the 21-centimeter line. In the context of quantum information, every time a 21-centimeter photon is emitted or absorbed by a hydrogen atom, a discrete quantum logic gate has been executed. The universe has effectively flipped a bit.

What makes this specific atomic qubit so crucial to the cosmic computational framework is its extreme stability. The transition from the parallel to the antiparallel state is highly forbidden by standard quantum selection rules. A solitary hydrogen atom in the parallel state left entirely undisturbed in a vacuum will take, on average, around 10 million years to spontaneously undergo a spin-flip transition. This phenomenally long half-life means that, absent external collisions, the neutral hydrogen atom is an incredibly stable memory register. It can hold a precise quantum state for millions of years without decohering, acting as a highly reliable, long-term physical storage medium for the local interstellar environment.

6.3 Nebular Hydrogen Clouds as Dilute Quantum Processors

While individual hydrogen atoms are exceptionally stable, the universe rarely leaves them undisturbed. The vast majority of the cosmos's hydrogen is not locked within the dense, fiery interiors of stars, but rather drifts through the dark, frigid expanses between them. These interstellar and intergalactic mediums are dominated by massive HI regions, which are colossal clouds of electrically neutral hydrogen spanning tens to hundreds of light-years across. Though they are categorized as clouds, these regions represent an extreme vacuum by terrestrial standards, typically containing only one to ten individual hydrogen atoms per cubic centimeter.

Despite this extreme dilution, the sheer volumetric scale of these nebulae transforms them into massive, localized data processing centers. Within these colossal clouds, hydrogen atoms slowly drift, occasionally colliding with one another, with stray cosmic rays, or with photons from distant starlight. Every collision has the potential to force a spin-flip transition, writing new data into the atomic qubit, or entangling the quantum states of the colliding particles. Therefore, an interstellar hydrogen cloud is not merely a static reservoir of gas; it is a continuously calculating quantum network.

The overall computational rate of these clouds is vastly slower than the energetic chaos of a star, but their slow, cold processing is precisely what dictates the large-scale evolution of galaxies. As these hydrogen atoms interact, flip their spins, and radiate away 21-centimeter photons, the entire cloud gradually loses thermal energy. This loss of heat eventually allows gravity to overcome the internal pressure of the gas, triggering the slow, methodical collapse of the cloud. The culmination of this gradual, localized quantum computation is the ignition of nuclear fusion and the birth of a new star. Thus, the massive, dilute arrays of atomic hydrogen qubits in the interstellar medium serve as the foundational processing hardware that computes the macroscopic physical architecture of the luminous universe.

7. Black Hole Thermodynamics and Entropy

Having traversed the dilute, sprawling data networks of cosmic background radiation and interstellar hydrogen, the architectural map of the universal quantum computer must now confront the absolute extremes of physical reality. If the vacuum of space acts as the universe's long-term memory and hydrogen clouds serve as its slow, distributed processors, black holes represent its ultimate, maximally compressed hard drives. For decades, classical general relativity treated black holes merely as gravitational sinks—regions of spacetime so violently warped that not even light could escape. However, the synthesis of general relativity, quantum field theory, and information theory has radically redefined the black hole. It is no longer viewed as a cosmic destroyer of matter, but rather as the universe's most efficient and dense repository of quantum information. To understand how a macroscopic gravitational singularity functions as a quantum storage device, one must examine the profound and unexpected mathematical equivalencies between the laws of thermodynamics and the mechanics of black holes. This section details the historical derivation of black hole entropy, explains the physical significance of the event horizon, and translates classical thermodynamic equations into discrete measurements of quantum information.

7.1 The Event Horizon and the Area Law

The boundary that defines a black hole is the event horizon, a theoretical threshold around the singularity where the escape velocity exactly equals the speed of light. In classical general relativity, the event horizon is a point of no return, a one-way membrane through which matter and radiation can enter but can never leave. According to the "no-hair theorem" of classical physics, once an object falls past this horizon, all of its complex, distinguishing characteristics are completely obliterated. The black hole retains only three observable macroscopic properties: total mass, electric charge, and angular momentum. From a classical perspective, a black hole formed by the collapse of a trillion tons of highly structured, complex biological

matter would be entirely indistinguishable from a black hole formed by the collapse of a trillion tons of random thermal photons.

However, this classical view introduced a massive paradox when viewed through the lens of thermodynamics. The Second Law of Thermodynamics dictates that the total entropy—the measure of disorder or hidden information—of an isolated physical system can never decrease over time. If a highly complex, high-entropy object is dropped into a black hole and its information is permanently erased from the observable universe, the total entropy of the universe would ostensibly decrease, violating one of the most sacrosanct laws of physics.

A critical breakthrough occurred when it was mathematically demonstrated that the total surface area of a black hole's event horizon can never decrease over time in any classical physical process. Even when two black holes collide and merge, the surface area of the resulting event horizon is always greater than or equal to the sum of the surface areas of the two original black holes. This strictly monotonic increase perfectly mirrored the relentless, one-way arrow of thermodynamic entropy. The mathematical behavior of the event horizon's area was entirely analogous to the behavior of entropy in a closed system, suggesting a profound, underlying physical connection between the geometry of spacetime and the laws of heat and disorder.

7.2 Deriving Bekenstein-Hawking Entropy

The realization that the event horizon's area behaves exactly like entropy led theoretical physicist Jacob Bekenstein to propose a radical hypothesis: the area of an event horizon does not merely resemble entropy, it fundamentally is entropy. Bekenstein argued that when matter falls into a black hole, its entropy is not destroyed; rather, it is added to the black hole, causing the event horizon to physically expand. This resolved the thermodynamic paradox by ensuring that the total entropy of the universe continues to increase.

This hypothesis was initially met with intense skepticism because, according to classical thermodynamics, anything that possesses entropy must also possess a physical temperature. If a black hole has a temperature greater than absolute zero, it must emit thermal radiation. But classical general relativity strictly forbade anything from escaping a black hole. This contradiction was famously resolved by Stephen Hawking, who applied quantum field theory to the curved spacetime immediately outside the event horizon. Hawking discovered that due to the constant creation and annihilation of virtual particle-antiparticle pairs in the quantum vacuum, a black hole does, in fact, emit a steady stream of thermal radiation. This phenomenon, known as Hawking radiation, proved that black holes have a measurable temperature and, consequently, a precise, calculable entropy.

The resulting mathematical formulation, known as the Bekenstein-Hawking entropy formula, stands as one of the most important equations in modern theoretical physics. It is written as S =

(k_B * A * c^3) / (4 * G * h-bar). In this equation, S represents the entropy of the black hole. The formula states that this entropy is directly proportional to the total surface area of the event horizon, represented by A. What makes this equation uniquely profound is its simultaneous incorporation of the fundamental constants from four vastly different realms of physics. It unifies thermodynamics via the Boltzmann constant (k_B), general relativity via the gravitational constant (G), special relativity via the speed of light (c), and quantum mechanics via the reduced Planck constant (h-bar). This single equation seamlessly binds the macroscopic geometry of gravity to the microscopic world of quantum information.

7.3 Translating Entropy into Qubits of Hidden Information

To fully integrate black holes into the framework of the cosmic quantum computer, thermodynamic entropy must be directly translated into the language of information theory. In statistical mechanics, entropy is a measure of the number of microscopic microstates that correspond to a single, observable macroscopic macrostate. In the context of a black hole, the macroscopic state is defined entirely by its mass, charge, and spin. The microscopic states are all the possible quantum arrangements of the matter and energy that collapsed to form the black hole in the first place. Therefore, the Bekenstein-Hawking entropy is essentially a definitive count of the hidden quantum information locked inside the black hole.

To quantify this hidden information, one must examine the denominator of the Bekenstein-Hawking formula, which consists of the Planck length squared. The Planck length, derived from combining the gravitational constant, the speed of light, and Planck's constant, represents the absolute smallest measurable scale of space in the universe. A Planck area is a two-dimensional square with sides equal to one Planck length. According to the Bekenstein-Hawking formula, the total entropy of a black hole is exactly equal to one quarter of the area of the event horizon when measured in Planck areas.

In terms of quantum computing, this means that the surface of a black hole's event horizon acts as a pixelated, spherical screen. Every four discrete Planck areas on this horizon encode exactly one fundamental unit of natural information, or one physical qubit. The sheer density of this information storage is unparalleled. A single square centimeter of a black hole's event horizon contains roughly 10^66 discrete Planck areas, meaning it stores an incomprehensible amount of quantum data in a microscopic amount of space. By translating thermodynamic entropy into a strict count of spatial pixels, physics demonstrates that a black hole is the absolute limit of data compression. It is a cosmic hard drive perfectly optimized by the laws of gravity, storing the maximum possible amount of quantum information physically permitted within a finite volume of spacetime.

8. Supermassive Black Holes: The Densest Information Repositories

While the mathematical framework of Bekenstein-Hawking entropy establishes that any black hole represents a state of maximum data compression, the absolute scale of this informational storage is wildly dependent on the physical size of the singularity in question. In the cosmos, black holes are not uniform; they exist across a vast spectrum of masses. Stellar-mass black holes, born from the collapse of massive stars, hold an immense amount of data, but they are entirely dwarfed by the gravitational behemoths that anchor the universe's largest structures. To locate the absolute densest and most massive localized repositories of quantum information in the cosmos, one must peer into the hearts of galaxies. Here, supermassive black holes do not merely participate in the cosmic computational network; they physically dominate its total memory capacity. This section explores the extreme mass concentrations of galactic centers, compares the staggering information density of these singularities against the dilute vacuum of space, and frames the famous black hole information paradox as a fundamental problem of cosmic data retrieval.

8.1 Galactic Centers and Extreme Mass Concentrations

Nearly every massive galaxy in the observable universe, including our own Milky Way, harbors a supermassive black hole at its rotational center. Unlike stellar-mass black holes, which typically range from five to a few hundred times the mass of our sun, supermassive black holes possess masses ranging from millions to tens of billions of solar masses. Their origins remain one of the most active areas of cosmological research, with theories suggesting they grew from the direct collapse of massive primordial gas clouds or the runaway mergers of dense star clusters in the early universe.

From an information-theoretic perspective, the defining characteristic of these galactic centers is not just their immense mass, but how that mass translates into thermodynamic entropy. For a standard, non-rotating Schwarzschild black hole, the radius of the event horizon is directly and linearly proportional to its mass. Because the surface area of a sphere scales with the square of its radius, the surface area of a black hole's event horizon—and therefore its total entropy and qubit capacity—scales with the square of its mass.

This mathematical relationship leads to a profound cosmic reality. If ten stellar-mass black holes merge together, the resulting black hole does not simply have ten times the information capacity; it has one hundred times the capacity. Because entropy scales quadratically with mass, supermassive black holes are disproportionately massive data vaults. A single supermassive black hole weighing one billion solar masses contains as much quantum information as one quintillion individual solar-mass black holes. Consequently, the universe's bound quantum

information is not evenly distributed among its gravitational wells; it is overwhelmingly monopolized by the singularities residing at the centers of galaxies.

8.2 Information Density Comparisons: Vacuum vs. Singularity

To truly grasp the architecture of the universal computer, one must compare the distributed memory of the vacuum with the concentrated memory of a supermassive black hole. As established in earlier sections, the Cosmic Microwave Background contains approximately 10^90 free-floating photonic qubits. However, this staggering amount of information is diluted across the entire observable volume of the universe, spanning roughly 3.5 x 10^86 cubic centimeters. The vacuum is a massively parallel but incredibly sparse data network.

In stark contrast, a single supermassive black hole, such as the one at the center of the giant elliptical galaxy M87, possesses an estimated mass of 6.5 billion solar masses. Calculating the Bekenstein-Hawking entropy for an object of this extreme mass yields an informational capacity of roughly 10^94 bits. This single localized object contains ten thousand times more quantum information than all the background radiation in the entire observable universe combined. Yet, the event horizon of the M87 black hole is only roughly the size of our solar system.

When physicists calculate the total entropy of the observable universe today, summing all the radiation, dust, planets, stars, and stellar remnants, the calculation is overwhelmingly dominated by supermassive black holes. The total qubit count of all supermassive black holes combined is estimated to be around 10^104. This means that 99.999 percent of the total localized, bound quantum information in the universe is sequestered on the two-dimensional surfaces of galactic centers. If the universe is a computer, the vacuum of space acts as a low-density, high-speed RAM carrying ancient cosmic data, while supermassive black holes serve as extreme-density, localized hard drives locking away the vast majority of the universe's total accrued entropy.

8.3 The Black Hole Information Paradox in a Computational Context

Treating a supermassive black hole as the ultimate cosmic hard drive inevitably leads to the most contentious debate in modern theoretical physics: the black hole information paradox. In a standard computational system, data written to a hard drive can theoretically be read, retrieved, or erased. However, the physical mechanics of black holes complicate this drastically. If a quantum system falls past the event horizon, its discrete information is added to the black hole's total entropy. But, as Stephen Hawking demonstrated, black holes are not entirely black; they emit Hawking radiation and slowly lose mass. Over unimaginable trillions of years, a supermassive black hole will completely evaporate into a cloud of thermal photons.

The paradox arises from the nature of this evaporation. According to Hawking's original calculations, this radiation is perfectly thermal and completely random. It carries no specific

information about the matter that initially fell into the black hole. If a black hole evaporates entirely into random noise, the highly structured quantum information of the matter that formed it is permanently destroyed. This permanently violates unitarity, a core tenet of quantum mechanics which dictates that quantum information can never be fundamentally erased from the universe.

In a computational context, the paradox asks whether a black hole is a permanent data sink that ultimately destroys information, or a highly chaotic quantum processor that eventually returns it. Modern resolutions to this paradox lean heavily on quantum information theory. Concepts such as the scrambling time dictate how fast a black hole can take incoming ordered qubits and disperse them across its event horizon via quantum entanglement. Advanced theoretical models suggest that as the black hole evaporates, the departing Hawking radiation is highly entangled with the internal states of the black hole. Eventually, past a threshold known as the Page time, the escaping radiation begins to carry the hidden quantum information back out into the universe. In this view, a supermassive black hole is not a cosmic trash can that permanently deletes data, but rather the universe's most extreme cryptographic scrambler, holding onto data for aeons before slowly leaking it back into the vast, expanding computational void.

9. The Holographic Principle and Universal Boundaries

The revelation that a supermassive black hole's total quantum information is entirely dictated by the two-dimensional surface area of its event horizon—rather than its three-dimensional volume—represents a profound crisis for classical spatial geometry. If the densest objects in the cosmos encode their internal states strictly on their boundaries, it forces a radical reevaluation of how information is distributed across the rest of the physical universe. This realization birthed the holographic principle, a sweeping theoretical framework suggesting that the area-scaling laws of black hole thermodynamics apply universally to all regions of spacetime. In the context of the cosmic quantum computer, the holographic principle provides the ultimate architectural blueprint. It defines the absolute upper limits of the universe's memory capacity and suggests that the three-dimensional processing of qubits we observe might be an emergent projection from a lower-dimensional boundary. This section explores the mechanics of encoding three-dimensional volume on two-dimensional surfaces, calculates the absolute maximum entropy bounds for the observable universe, and examines the profound implications this holographic nature has for the true spatial distribution of cosmic data.

9.1 Encoding 3D Volume on a 2D Boundary

In our standard, intuitive experience of classical physics, the capacity of any container to hold physical objects or information scales with its three-dimensional volume. If you double the

radius of a spherical hard drive, its volume—and theoretically its data capacity—increases by a factor of eight. However, the Bekenstein-Hawking entropy formula destroyed this assumption for gravitational extremes. If a region of space accumulates too much mass and information, it collapses into a black hole, and its maximum information capacity suddenly scales exclusively with the square of its radius, representing the two-dimensional surface area.

Pioneered by theoretical physicists Gerard 't Hooft and Leonard Susskind, the holographic principle generalizes this black hole phenomenon to encompass all of physical reality. The principle asserts that the complete physical description of any three-dimensional volume of space can be fully mathematically encoded on the two-dimensional boundary enveloping that region. Much like an optical hologram uses a flat, two-dimensional plate of interference patterns to project a fully three-dimensional image, the universe may be structured such that the three-dimensional phenomena we observe—particles, forces, and galactic structures—are emergent projections of quantum data residing on a distant, lower-dimensional bounding surface.

In this paradigm, the fundamental pixels of reality are not three-dimensional cubic blocks of volume, but two-dimensional Planck areas residing on a cosmic boundary. Each of these Planck areas can hold exactly one binary unit of information, or one qubit. This strict dimensional reduction drastically limits the amount of independent degrees of freedom available to a region of space. It mathematically dictates that if you attempt to pack a volume of space with independently oscillating, maximally dense qubits, you will reach a hard physical limit—the holographic bound—long before the volume appears classically "full." Once this bound is reached, packing even a single additional qubit into the region will inevitably trigger gravitational collapse, transforming the localized computing node into a black hole.

9.2 Maximum Entropy Bounds for the Observable Universe

Applying the holographic principle to the universe at large allows physicists to calculate the absolute maximum memory capacity of the entire cosmic computer. To do this, we must define the ultimate boundary of our localized reality: the cosmological event horizon. Because the universe has been expanding for 13.8 billion years, and because the speed of light is finite, there is a spherical boundary beyond which we can never observe or interact with any physical phenomena. This cosmological horizon envelops the observable universe, acting as the outermost screen of our physical reality.

By applying the holographic bound to the surface area of this cosmic horizon, we can determine the maximum number of qubits the observable universe could possibly contain without collapsing into a singular universe-spanning black hole. The current comoving radius of the observable universe is approximately 46.5 billion light-years. Calculating the surface area of a

sphere with this radius, and dividing that area into discrete, microscopic Planck areas, yields an astonishing number: roughly 10^122 Planck areas.

Because the holographic principle equates one quarter of these discrete areas to one qubit of information, the absolute maximum computational capacity of the observable universe is approximately 10^122 qubits. This colossal number represents the theoretical ceiling of universal computation. It is the ultimate limit of how much data can be written into the fabric of spacetime. Crucially, when we compare this absolute ceiling to the actual known contents of the universe, a profound discrepancy emerges. As established in previous sections, the total free-floating radiation of the Cosmic Microwave Background accounts for roughly 10^90 qubits, and the extreme concentrations of supermassive black holes account for roughly 10^104 qubits. Even when accounting for every known particle and singularity, the universe is operating immensely far below its maximum holographic capacity. The universe is not a full hard drive; it is an incredibly vast, mostly empty computational grid executing algorithms in a tiny fraction of its available memory space.

9.3 Implications for the Spatial Distribution of Qubits

If the holographic principle is taken to its logical conclusion, it fundamentally dismantles our understanding of where quantum information actually resides. When we map the universe, we instinctively place the photonic qubits of the Cosmic Microwave Background traversing the vast 3D void, and we place the massive qubit vaults of black holes residing at the 3D centers of galaxies. However, if the entire 3D volume of the observable cosmos is merely an emergent projection of a 2D boundary, then the true, fundamental location of all these qubits is not the interior volume of space at all, but rather the cosmological horizon itself.

This framework implies that spatial distribution is an illusion generated by quantum entanglement. The apparent vast distances between a hydrogen qubit in the Milky Way and a relic neutrino traversing the Andromeda galaxy might simply correspond to how strongly their respective data points are entangled on the 2D boundary. Highly entangled qubits on the boundary project mathematically as being spatially close in the 3D interior, while unentangled data points project as being separated by vast cosmic voids.

Consequently, the architecture of the universal quantum computer may not resemble a classical 3D network of wires and nodes. Instead, it may function as a closed, 2D quantum processing unit mapping its internal algorithms into the illusion of three-dimensional depth. The cosmic microwave photons, the neutral hydrogen clouds, and the supermassive black holes are not physically separate objects drifting through a vast container; rather, they are distinct algorithms and localized data clusters running simultaneously on the two-dimensional cosmic screen. This holographic perspective perfectly bridges the gap between quantum mechanics and general

relativity, suggesting that gravity and three-dimensional space itself are not fundamental features of the cosmic hardware, but merely the output display of an underlying, lower-dimensional quantum computation.

10. Dark Matter and Dark Energy: Unknowns in Cosmic Computation

While the preceding sections have meticulously mapped the quantum information residing in photons, neutrinos, baryonic matter, and black holes, this accounting covers only the luminous and interactive fraction of the cosmos. According to the standard model of cosmology, standard baryonic matter and radiation constitute less than five percent of the universe's total mass-energy budget. The remaining ninety-five percent is entirely opaque to electromagnetic observation, divided into two mysterious components: dark matter, which accounts for roughly twenty-seven percent, and dark energy, which dominates with roughly sixty-eight percent. If the universe operates as a holistic quantum computer, this massive dark sector cannot simply be ignored. It must play a fundamental, structural role in the cosmic architecture, either as a vast reservoir of hidden qubits or as the underlying physical mechanism driving the expansion of the computational grid itself. This section attempts to integrate these invisible components into the information-theoretic paradigm by exploring the potential qubit capacity of theoretical dark matter particles, analyzing how dark energy perpetually alters the physical boundaries of universal computation, and examining leading theories that model the dark sector purely as emergent phenomena of quantum information.

10.1 Potential Information Carrying Capacity of Dark Matter Candidates

Dark matter is strictly defined by its gravitational influence on visible structures. It forms the massive invisible halos that envelop galaxies, provides the gravitational scaffolding for the cosmic web, and dictates the rotational velocities of stars. Because it does not interact with the electromagnetic force, it neither emits nor absorbs light, making its exact particulate nature one of the greatest unresolved mysteries in modern physics. However, if dark matter is composed of fundamental particles, as is widely assumed, then by the laws of quantum mechanics, these particles must possess intrinsic degrees of freedom. Therefore, they must act as carriers of quantum information.

The potential information capacity of dark matter depends entirely on its theoretical composition. If dark matter consists of Weakly Interacting Massive Particles, these heavy, slow-moving fermions would possess internal spin states. Due to their high mass and extremely low interaction cross-sections, they would function as highly stable, localized qubits. Unlike the relativistic neutrinos of the cosmic background, these massive particles would be gravitationally bound within galactic halos, forming cold, dense, invisible memory banks. Because they rarely

collide or interact via anything other than gravity and the weak nuclear force, their quantum states would experience almost zero decoherence.

Alternatively, if dark matter is composed of axions—hypothetical, ultra-lightweight bosons—the informational architecture changes dramatically. Axions are theorized to exist in a Bose-Einstein condensate state across galactic scales, meaning massive numbers of them would occupy the exact same macroscopic quantum state. In this scenario, a dark matter halo would not act as a collection of discrete, independent qubits, but rather as a single, coherent, macroscopic quantum object. Regardless of the specific particle, calculating the total thermodynamic entropy of the dark matter sector suggests it holds significantly more information than the cosmic microwave background, yet vastly less than supermassive black holes. Dark matter likely functions as the universe's most robust, decoherence-free structural memory, silently shaping the physical processors of baryonic matter without directly participating in their chaotic algorithms.

10.2 Dark Energy and the Expanding Computational Space

While dark matter acts as the gravitational glue holding localized computing nodes together, dark energy acts as a relentless, repulsive force driving them apart. Discovered through the observation of distant supernovae, dark energy is responsible for the accelerating expansion of the universe. In standard cosmology, it is often mathematically modeled as the cosmological constant—an intrinsic, irreducible energy density inherent to the vacuum of space itself. In the framework of quantum information, dark energy presents a profound complication to the cosmic computer by continuously and aggressively altering the physical dimensions of the processing space.

As dark energy accelerates cosmic expansion, the distances between gravitationally unbound galaxy clusters increase faster than the speed of light. This means that qubits separated by vast intergalactic distances can no longer exchange information. The universal computer is actively fragmenting. While the overall space of the universe expands, the observable universe—the region from which light can currently reach an observer—is surrounded by a cosmological event horizon. Because of dark energy, this horizon is shrinking relative to the expanding matter. Galaxies that were once part of our observable computational network are crossing this horizon and being permanently deleted from our local cosmic dataset.

However, there is a counter-intuitive informational aspect to this expansion. Just as a black hole has an event horizon with a measurable Bekenstein-Hawking entropy, the cosmological horizon of an accelerating universe, known as de Sitter space, also possesses an intrinsic thermodynamic entropy. As dark energy dominates the cosmos, driving it toward an empty, cold state, the total entropy of the universe does not approach zero. Instead, the entropy of the

vacuum itself maximizes. The accelerating expansion effectively writes information onto the outward boundary of the observable universe. In the far future, when all stars have died and all black holes have evaporated, the final, maximum computational state of the universe will be governed entirely by the entropy of the dark energy vacuum, a static, maximum-capacity hard drive with no remaining matter to process its data.

10.3 Theoretical Models for Dark Qubits

The struggle to identify the physical particles of dark matter and the exact energy field of dark energy has led some theoretical physicists to propose that these phenomena might not be distinct physical substances at all. Instead, they might be emergent properties of quantum information processing itself. This radical approach flips the standard model: rather than particles serving as the hardware for quantum information, quantum information serves as the fundamental bedrock from which gravity, space, and dark phenomena emerge.

One leading model in this domain is the theory of emergent or entropic gravity. In this framework, gravity is not a fundamental force mediated by gravitons, but rather a macroscopic thermodynamic consequence of quantum entanglement between discrete qubits on a holographic boundary. When physicists mathematically model how the entanglement entropy of the quantum vacuum behaves on large scales, standard Newtonian and Einsteinian gravity emerge naturally. Crucially, when these models are pushed to galactic scales where the acceleration falls below a specific threshold, the standard laws of gravity break down. The resulting mathematical corrections perfectly mimic the exact gravitational effects attributed to dark matter halos. In this purely information-theoretic model, dark matter does not consist of hidden particles; it is simply the thermodynamic signature of the quantum vacuum's entanglement structure becoming highly pronounced on galactic scales.

Similarly, dark energy can be interpreted through the lens of quantum entanglement. The irreducible energy of the vacuum may represent the baseline entanglement entropy of the spacetime fabric. As the universe computes, the continuous entanglement of primordial qubits generates an outward pressure. In these theoretical models, the dark sector is entirely composed of dark qubits—not unknown particles, but pure, abstract nodes of quantum information whose vast network of complex correlations generates the geometric illusions we perceive as missing mass and expanding space. By translating the dark sector into the language of qubits and entanglement entropy, quantum information cosmology offers a potential pathway to resolving the two greatest physical anomalies of the universe without requiring the invention of new, undetectable elementary particles.

11. Cosmological Quantum Entanglement

The preceding sections have detailed the physical hardware of the universe, cataloging the discrete physical qubits stored within background radiation, hydrogen clouds, and the extreme geometry of black holes. However, a computer consisting merely of isolated memory registers is incapable of executing complex algorithms. For computation to occur, qubits must interact, share information, and become deeply correlated. In the realm of quantum mechanics, this operational linkage is achieved through quantum entanglement. Entanglement is the physical phenomenon whereby multiple particles become linked such that the quantum state of one particle cannot be described independently of the state of the others, regardless of the spatial distance separating them. If physical particles are the memory registers of the cosmos, entanglement serves as the invisible wiring that connects them, facilitating the massively parallel processing that defines quantum computation. To understand the universe as a cohesive computational entity, one must trace the history of this invisible wiring. This section explores how the rapid expansion of the early universe stretched microscopic quantum correlations to cosmic scales, examines theoretical proposals for detecting these primordial Bell correlations in modern astronomical data, and explains how the continuous loss of entanglement drives the emergence of the classical physical reality we observe today.

11.1 Primordial Entanglement and Cosmic Inflation

The story of the universe's interconnected wiring begins in the first fraction of a second following the Big Bang, during an epoch known as cosmic inflation. According to standard cosmological models, at approximately 10^-36 seconds after the initial singularity, the universe underwent an incredibly violent, exponential expansion. In a mere fraction of a second, the spatial volume of the universe increased by a factor of at least 10^78. Before this inflationary period, the entirety of the observable universe was compressed into an incredibly hot, dense, and microscopic quantum patch. Because this patch was so unimaginably compact, all the fundamental fields and primordial particles within it were in constant contact, engaging in continuous quantum interactions that resulted in a state of maximum, universe-wide quantum entanglement.

When inflation triggered the exponential stretching of space, it fundamentally altered the spatial relationship of these entangled quantum states. Quantum fluctuations—microscopic, temporary changes in the amount of energy in a point in space—were rapidly stretched from the subatomic scale to macroscopic, cosmological scales. Crucially, the mathematical framework of quantum mechanics dictates that stretching space does not sever quantum entanglement. If two virtual particles were perfectly entangled in the pre-inflationary vacuum, the violent expansion of the universe simply pulled them apart, placing them at opposite ends of the cosmos while preserving their shared quantum state. Therefore, cosmic inflation acted as an

ultimate distribution mechanism for the cosmic computer's wiring. It generated a pervasive, underlying web of long-range quantum correlations, ensuring that the macroscopic large-scale structure of the universe was fundamentally seeded by microscopic entangled quantum fluctuations.

11.2 Bell Correlations across the Cosmic Microwave Background

If cosmic inflation truly stretched entangled pairs of qubits across intergalactic distances, the observable universe should still harbor the mathematical signatures of this primordial quantum wiring. In laboratory settings, physicists prove the existence of entanglement between two particles by testing for violations of Bell inequalities. A Bell inequality is a mathematical theorem demonstrating that if two particles act entirely independently, their statistical correlations can only reach a certain maximum limit. When laboratory particles violate this limit, it definitively proves they are quantum mechanically entangled and acting as a single, unified system.

In the context of quantum information cosmology, theoretical physicists are currently exploring whether it is possible to observe violations of Bell inequalities across the vast expanse of the cosmic microwave background. The temperature fluctuations and polarization patterns mapped onto the cosmic microwave background are direct physical imprints of the primordial quantum fluctuations stretched by inflation. If the universe is a fully integrated quantum computer, the polarization state of a photon arriving from the extreme northern hemisphere of the observable universe should theoretically exhibit subtle, non-classical statistical correlations with a photon arriving from the extreme southern hemisphere, despite the fact that these two regions of space have been separated by billions of light-years and have never been in classical causal contact.

While detecting these cosmic Bell correlations is exceptionally difficult due to the sheer noise and complexity of astrophysical data, advanced mathematical models suggest that the polarization patterns of relic photons carry specific parity signatures that cannot be explained by classical physics. By analyzing the massive datasets collected by modern orbital observatories, researchers are searching for specific statistical anomalies in the cosmic microwave background that would prove different regions of the sky are deeply entangled. The successful detection of a cosmological Bell inequality violation would provide the ultimate, indisputable proof that the universe operates as a single, interconnected quantum processing network, wired together in the first moments of time.

11.3 Decoherence and the Classical Transition of the Universe

If the early universe was a state of maximum quantum entanglement, a glaring physical question remains: why does the universe we inhabit today appear so strictly classical? In our daily macroscopic experience, objects do not exist in superpositions of multiple states, and we

do not observe quantum entanglement between baseballs or planets. In the framework of quantum computation, this transition from a purely quantum state to a classical reality is driven by a phenomenon known as quantum decoherence.

Decoherence is the process by which a quantum system loses its fragile, complex quantum superpositions due to continuous interaction with its surrounding environment. In a human-engineered quantum computer, a stray photon or a slight fluctuation in magnetic fields can interact with a qubit, causing its wave function to collapse and destroying the quantum calculation. In the cosmological computer, the environment is the universe itself. As the universe expanded and cooled, particles began to interact, collide, and bind together to form complex structures. Every time a primordial cosmic microwave background photon scattered off a free electron, or every time a hydrogen atom underwent a spin-flip transition, the environment essentially performed a measurement on that quantum state.

This continuous, relentless barrage of physical interactions caused the vast majority of the universe's primordial superpositions to collapse. As quantum information leaked out into the broader environment, the pristine entanglement of the early universe was gradually scrambled into localized thermal noise. This process, often described by theoretical physicists as quantum Darwinism, allows only the most stable, redundant physical states to survive the chaotic environment of the cosmos. The classical world of planets, stars, and galaxies is not a separate physical realm from the quantum world; rather, it is the emergent, localized output of a quantum computer that has undergone relentless, universe-wide decoherence. While the fundamental hardware of the cosmos remains entirely quantum mechanical, the continuous interaction of its innumerable qubits creates the robust, classical illusion we experience as physical reality.

12. Future Prospects in Observational Quantum Cosmology

For decades, the proposition that the universe functions as a colossal quantum computer remained a largely philosophical interpretation of theoretical physics. However, the rapidly advancing capabilities of modern astrophysics, combined with the exponential growth in human-engineered quantum technology, are actively driving this paradigm out of the realm of pure theory and into the domain of observational science. The ultimate goal of quantum information cosmology is to directly measure, map, and decode the quantum state of the universe using empirical data. As observational instruments achieve unprecedented levels of sensitivity, physicists are developing rigorous methodologies to detect the subtle, macroscopic signatures of cosmic quantum information processing. This final section outlines the most promising observational techniques currently being developed to measure cosmological qubits,

explores how terrestrial quantum computers will be used to simulate and decode astrophysical datasets, and provides a definitive conclusion on the overarching computational architecture of the cosmos.

12.1 Proposed Detection Methods for Cosmological Qubit Signatures

To empirically validate the cosmic computational network, astrophysicists must develop instruments capable of reading the discrete states of the universe's most abundant qubits. The most immediate prospect lies in the precise measurement of the Cosmic Microwave Background. While the temperature fluctuations of this primordial radiation have been mapped with exquisite precision, the next frontier is high-resolution polarimetry. Specifically, researchers are searching for B-mode polarization patterns—a distinct twisting in the alignment of the photonic qubits. These primordial B-modes are theorized to have been generated by gravitational waves during cosmic inflation. If these specific polarization signatures are detected, they would serve as a direct observational record of the entangled quantum fluctuations that occurred a fraction of a second after the Big Bang, essentially providing a snapshot of the universe's initial algorithmic input.

Beyond photons, detecting the fermionic qubits of the Cosmic Neutrino Background represents one of the greatest technical challenges in modern experimental physics. Because relic neutrinos possess incredibly low energy and almost never interact with baryonic matter, they cannot be observed using conventional telescopes. However, next-generation experimental frameworks propose using large quantities of radioactive isotopes, such as tritium, which undergo beta decay. The theoretical premise is that a collision with a relic neutrino could induce a specific, measurable shift in the energy spectrum of the emitted electrons. Successfully capturing a relic neutrino would allow physicists to directly measure the oldest, most isolated qubits in existence, providing pristine data regarding the weak force interactions that dominated the first second of universal computation.

Finally, observing the active, baryonic processing network relies heavily on the advancement of massive radio telescope arrays. These arrays are being constructed to map the 21-centimeter emission line of neutral hydrogen across billions of light-years, specifically targeting the cosmic dark ages and the epoch of reionization. By charting exactly where and when neutral hydrogen clouds underwent spin-flip transitions, astrophysicists are essentially tracking the execution of atomic logic gates on a cosmological scale. Mapping this data in three dimensions will reveal how the slow, dilute computation of the interstellar medium ultimately organized the first chaotic clouds of gas into the structured, luminous galaxies we observe today.

12.2 Integrating Quantum Computing with Astrophysical Data

The effort to decode the universe's computational architecture will not rely solely on telescopes; it will profoundly depend on the application of human-engineered quantum processors. As terrestrial quantum computers scale from dozens of noisy qubits to thousands of fault-tolerant logical qubits, they will become indispensable tools for simulating the extreme physics of the cosmos. Classical supercomputers are fundamentally incapable of efficiently simulating the quantum entanglement and non-linear dynamics inherent in black hole thermodynamics or the chaotic plasma of the early universe. By mapping the Hamiltonian equations of early universe cosmology directly onto the architecture of a laboratory quantum processor, physicists can effectively run small-scale simulations of cosmic inflation and primordial entanglement that classical physics cannot compute.

Furthermore, laboratory quantum computers are actively being used to test the holographic principle and black hole information dynamics. Theoretical physicists have already begun running quantum algorithms, such as the Sachdev-Ye-Kitaev model, which mathematically behave like simplified, one-dimensional black holes. By purposefully scrambling information inside a quantum processor and measuring how it is distributed among the system's qubits, researchers can empirically study the exact mechanisms of Hawking radiation and quantum teleportation in a controlled laboratory setting. These localized quantum experiments provide direct analogs for the information processing occurring on the event horizons of supermassive black holes located millions of light-years away.

Additionally, quantum machine learning algorithms will eventually be deployed to analyze the staggeringly massive datasets generated by next-generation cosmological surveys. Standard classical algorithms struggle to identify subtle, non-local correlations hidden within the petabytes of data produced by cosmic microwave background maps and deep-sky galaxy surveys. Quantum machine learning, operating on the principles of quantum superposition and interference, is theoretically optimized to sift through this noise and detect the faint statistical anomalies indicative of cosmological Bell inequality violations or topological defects in the fabric of spacetime. The union of artificial quantum computing and observational astrophysics will ultimately provide the mathematical translation key needed to read the universe's underlying code.

12.3 Final Conclusions on the Universe's Computational Architecture

The synthesis of quantum mechanics, general relativity, and information theory inevitably leads to a single, profound conclusion: the universe is not a static collection of inert matter and empty space, but a dynamic, self-executing physical algorithm. Every fundamental particle represents a physical register of information, and every interaction between those particles is a discrete

logical operation driving the system forward in time. This paper has comprehensively mapped the architecture of this cosmic computer. The foundational memory capacity of the universe is distributed across the vacuum of space, composed of approximately 10^90 free-floating photonic qubits in the Cosmic Microwave Background and an equally massive array of oscillating fermionic qubits in the Cosmic Neutrino Background. These relic particles serve as the ultimate long-term data storage, carrying the boundary conditions of the early universe across the expanse of time with virtually zero decoherence.

Conversely, the active processing of this quantum data occurs within gravitationally bound systems. The dilute clouds of neutral hydrogen permeating the interstellar medium act as the universe's default atomic logic gates, slowly computing the large-scale thermodynamic evolution of galactic structures through discrete spin-flip transitions. At the absolute extremes of physical reality, supermassive black holes function as the universe's ultimate data compressors. Governed by the Bekenstein-Hawking entropy formula and the holographic principle, these singularities sequester the vast majority of the universe's localized, bound quantum information upon their two-dimensional event horizons, transforming the chaotic thermodynamic collapse of matter into perfectly ordered, highly encrypted data storage.

Ultimately, the information-theoretic paradigm shifts our understanding of reality at its deepest level. It demonstrates that the thermodynamic clock of the universe is driven entirely by the processing of quantum states, bounded rigidly by the speed of light, the Planck length, and the continuous acceleration of dark energy. While the vast darkness of the cosmos may appear chaotic and empty, it is, in fact, relentlessly calculating. From the primordial quantum entanglement of the Big Bang to the slow, eventual evaporation of the final black holes, the entire history of the cosmos is the history of a single, finite quantum computation unfolding across the canvas of spacetime.

13. References

Bekenstein, J. D. (1973). Black holes and entropy. Physical Review D, 7(8), 2333-2346. https://doi.org/10.1103/PhysRevD.7.2333

Bremermann, H. J. (1962). Optimization through evolution and recombination. In M. C. Yovits, G. T. Jacobi, & G. D. Goldstein (Eds.), Self-Organizing Systems (pp. 93-106). Spartan Books.

Hawking, S. W. (1975). Particle creation by black holes. Communications in Mathematical Physics, 43(3), 199-220. https://doi.org/10.1007/BF02345020

Lloyd, S. (2000). Ultimate physical limits to computation. Nature, 406(6799), 1047-1054. https://doi.org/10.1038/35023282

Lloyd, S. (2002). Computational capacity of the universe. Physical Review Letters, 88(23), 237901. https://doi.org/10.1103/PhysRevLett.88.237901

Maldacena, J. (1999). The large N limit of superconformal field theories and supergravity. International Journal of Theoretical Physics, 38(4), 1113-1133. https://doi.org/10.1023/A:1026654312961

Margolus, N., & Levitin, L. B. (1998). The maximum speed of dynamical evolution. Physica D: Nonlinear Phenomena, 120(1-2), 188-195. https://doi.org/10.1016/S0167-2789(98)00054-2

Page, D. N. (1993). Information in black hole radiation. Physical Review Letters, 71(23), 3743-3746. https://doi.org/10.1103/PhysRevLett.71.3743

Susskind, L. (1995). The world as a hologram. Journal of Mathematical Physics, 36(11), 6377-6396. https://doi.org/10.1063/1.531249

't Hooft, G. (1993). Dimensional reduction in quantum gravity. arXiv preprint gr-qc/9310026. https://arxiv.org/abs/gr-qc/9310026

Verlinde, E. (2011). On the origin of gravity and the laws of Newton. Journal of High Energy Physics, 2011(4), 29. https://doi.org/10.1007/JHEP04(2011)029

Wheeler, J. A. (1990). Information, physics, quantum: The search for links. In W. Zurek (Ed.), Complexity, Entropy, and the Physics of Information (pp. 3-28). Addison-Wesley.

The author has published over 5,400 AI-assisted papers at: https://www.researchgate.net/profile/Douglas-Youvan/research . Google Scholar has indexed these preprints, and if you want a particular reference, the AI mode of a Google search (Gemini) has efficiently catalogued these preprints.